Draft version August 10, 2012
Preprint typeset using LaTeX style emulateapj v. 5/2/11

arXiv:1208.1761v1 [astro-ph.CO] 8 Aug 2012

RELAXATION OF BLAZAR INDUCED PAIR BEAMS IN COSMIC VOIDS

Francesco Miniati
Physics Department, Wolfgang-Pauli-Strasse 27, ETH-Zürich, CH-8093, Zürich, Switzerland; fm@phys.ethz.ch

Andrii Elyiv
Institut d'Astrophysique et de Géophysique, Université de Liège, 4000 Liège, Belgium and
Main Astronomical Observatory, Academy of Sciences of Ukraine, 27 Akademika Zabolotnoho St., 03680 Kyiv, Ukraine

Draft version August 10, 2012

ABSTRACT

The stability properties of a low density ultra relativistic pair beam produced in the intergalactic medium by multi-TeV gamma-ray photons from blazars are analyzed. The problem is relevant for probes of magnetic field in cosmic voids through gamma-ray observations. In addition, dissipation of such beams could affect considerably the thermal history of the intergalactic medium and structure formation. We use a Monte Carlo method to quantify the shower properties, in particular the bulk Lorentz factor and the angular spread of the beam, as a function of distance from the blazar. We find that the fastest growing modes, like any perturbation mode with even a very modest component perpendicular to the beam direction, requires a kinetic treatment. Combined with the effect of non-linear Landau damping, which suppresses the growth of plasma oscillations, the beam relaxation timescale is found typically longer than a Hubble time. Finally, density inhomogeneities associated with cosmic structure induce severe loss of resonance between the beam particles and plasma oscillations, strongly inhibiting their growth. We conclude that relativistic pair beams produced by blazars in the intergalactic medium are stable on timescales much longer compared to the electromagnetic cascade's. There appears to be no effect of pair-beams on the intergalactic medium.

Subject headings: gamma rays: general – instabilities – intergalactic medium – plasmas – radiation mechanisms: non-thermal – relativistic processes

1. INTRODUCTION

Streaming relativistic particles are common in tenuous astrophysical plasma and their propagation and stability properties a recurrent theme. Examples include transient hard X-ray emission in the solar corona (Benz 1993), quasars' jets (Lesch & Schlickeiser 1987), cosmic-rays streaming out of star forming galaxies (Miniati & Bell 2011) and cosmic-ray transport in the intracluster medium (Ensslin et al. 2011). Propagation and stability properties, related in particular to the exitation of plasma waves, is subject of attentive investigation as they can play a crucial role in the interpretation of observational data.

Ultra-relativistic beams of $e^{\pm}$ pairs are also generated in the intergalactic medium (IGM) by very high energy gamma-rays from distant blazars, by way of photon-photon interactions with the extragalactic background light (EBL, Gould & Schréder 1967; Schlickeiser et al. 2012). While blazars' spectra, and in particular their multi-TeV cut-off features, have been studied in detail to constraint the EBL (e.g. Aharonian et al. 2006), recently multi-GeV and TeV blazars observations have also been used to constrain magnetic field in cosmic voids for the first time (Neronov & Vovk 2010; Tavecchio et al. 2010; Dermer et al. 2011; Taylor et al. 2011). In fact, for flat enough blazar's spectra, the electromagnetic cascade should produce an observable spectral bump at multi-GeV energies. The absence of such a feature in a number of observed blazars is ascribed to the presence of a sufficiently strong magnetic field, $B_{\rm v} \gtrsim 10^{-18}$G, to deflect the pairs in less then an inverse Compton length, $\ell_{IC} \simeq {\rm Mpc}\,(E_{\pm}/{\rm TeV})^{-1}(1+z)^{-4}$, where z is the cosmological redshift (Plaga 1995; Neronov & Semikoz 2009).

However, in principle the pair-beam is subject to various instabilities, in particular microscopic plasma instabilities of the two-stream family. On this account, Broderick et al. (2012) conclude that transverse modes of the two-stream instability act on much shorter timescales than inverse Compton scattering, effectively inhibiting the cascade and invalidating the above magnetic field measurements. In addition, as a result of the beam's relaxation, substantial amount of energy would be deposited into the IGM, with dramatic consequences for its thermal history (Chang et al. 2012; Pfrommer et al. 2012).

In this paper, we reanalyze the stability of blazars induced ultra-relativistic pair beams. In particular, we use a Monte Carlo model of the electromagnetic shower to quantify the beam properties at various distances from the blazar, and analyze the stability of the produced beam following the work of Breizman, Rytov and collaborators (reviewed in Breizman & Ryutov 1974; Breizman 1990). We pay particular attention to the role of small, but finite, angular spread of the beam. We find that even for very modest perpendicular components of the wave-vector, the analysis of the instability requires a kinetic treatment. In addition, when non-linear Landau damping is taken into account, the growth rate of plasma oscillations is suppressed, so that the beam relaxation timescale becomes longer than a Hubble time. Finally, the resonance condition cannot be maintained in the presence of density inhomogeneities associated to cosmological structure formation, which also act to dramatically suppress the beam instability. Non-linear Landau damping was not explicitly included in Broderick's calculations and density inhomogeneities not considered, but they both appear to considerably stabilize the beam during a Hubble time, thereby ruling out effects of blazars' beam on the thermal history of the IGM and structure formation.

The rest of this paper is organized as follows. Sec. 2 summarizes the physical properties of pair beams produced by blazars and present the results of the Monte Carlo model. The two-stream instability in both the reactive and kinetic regimes is discussed in Sec. 3. Nonlinear effects are discussed in Sec. 4, where the timescales for the relaxation of the pair beam are also derived. Finally, Sec. 5 briefly summarizes the results.

2. PAIR BEAMS IN VOIDS

The unstable behavior of an ultra-relativistic pair beam depends, amongst others, on its density contrast to the IGM, Lorentz factor, angular and velocity spread. These characteristic quantities derive from the energy and number density of the pair producing photons, i.e. the blazar's spectral flux, F_γ, and the EBL model. The process has been studied extensively in the literature (e.g. Gould & Schréder 1967; Bonometto & Rees 1971; Schlickeiser et al. 2012) and in the following we briefly summarize its qualitative features before we describe the results of our Monte Carlo model of a blazar induced cascade.

2.1. *Basic Qualitative Features*

Pairs are most efficiently created just above the energy threshold for production, i.e. where

$$s \equiv E_\gamma E_{\rm EBL}(1-\cos\phi)/2m_e^2c^4 \geq 1, \tag{1}$$

with ϕ the angle between the interacting photons, E_γ and $E_{\rm EBL}$ the energy of the incident and target EBL photon, respectively, and the relativistic invariant, s, the center of mass energy square in units $m_e^2c^4$. The mean free path for the process depends on the details of the EBL model (Kneiske et al. 2004; Franceschini et al. 2008) but is approximately

$$\ell_{\gamma\gamma} \simeq 0.8 \left(\frac{E_\gamma}{\rm TeV}\right)^{-1} (1+z)^{-\zeta}{\rm Gpc}, \tag{2}$$

with $\zeta = 4.5$ for $z \leq 1$, $\zeta = 0$ otherwise (Neronov & Semikoz 2009; Broderick et al. 2012). The particle number density of the beam, n_b, is set by the balance of pair production rate, $2F_\gamma/\ell_{\gamma\gamma}$, evaluated close to production threshold, and energy loss rate. If inverse Compton losses dominate then, at a distance D from the blazar such that $F_\gamma = L_\gamma/4\pi D^2$,

$$\begin{aligned} n_b \simeq & \,2\frac{F_\gamma}{c}\frac{\ell_{IC}}{\ell_{\gamma\gamma}} \simeq 3\times10^{-25}{\rm cm}^{-3}\left(\frac{E_\gamma L_\gamma}{10^{45}{\rm erg/s}}\right) \\ & \times\left(\frac{D}{\rm Gpc}\right)^{-2}\left(\frac{E_\gamma}{\rm TeV}\right)^{-1}(1+z)^{\zeta-4}. \end{aligned} \tag{3}$$

where, $E_\gamma L_\gamma$, is an estimate of the blazar's equivalent isotropic gamma-ray luminosity for a source at distance D (see, Broderick et al. 2012). Each pair particle carries about half the energy of the incident gamma-ray, so the beam Lorentz factor is



FIG. 1.— *Top*: Peak (dash) and median (solid) pair beam energy as a function of distance from the blazar. Vertical bars correspond to 68% percentile of the energy spread about the median. A power law, $dn_\gamma/dE_\gamma \propto E_\gamma^{-2}$, in the range $10^3 \leq E_\gamma/m_ec^2 \leq 10^8$, is assumed for the blazar's spectrum, together with the *nominal* EBL model from Aharonian (2001). *Bottom*: Angular distribution of beam pairs generated at 55 Mpc (solid), 350 Mpc (dash) and 1 Gpc (dot-dash) from the blazar. The colored curves indicate the contribution to the angular distribution for 350 Mpc from pairs in the energy range: 10^4–10^5 (red), 10^5–10^6 (yellow), 10^6–10^7 (green), 10^7–10^8 (blue), respectively.

$$\Gamma_{beam} = \frac{E_\gamma}{2m_ec^2} \sim 10^6 \left(\frac{E_\gamma}{\rm TeV}\right)^{-1}. \tag{4}$$

Another important characteristic quantity of the beam is its angular spread, $\Delta\theta_{beam}$, determined by the distribution of angles θ between of pair produced particles with respect to the parent photon direction. This can be found to be related to Γ_{beam} and the relativistic invariant as

$$\Delta\theta_{beam} \leq \frac{s}{\Gamma_{beam}}\left(1-\frac{1}{s}\right)^{\frac{1}{2}}. \tag{5}$$

2.2. *Monte Carlo Model*

The characteristic quantities of a pair beam generated by gamma-rays from a blazar were computed using a Monte Carlo model of the electromagnetic cascade. The Monte Carlo code is fully described in Elyiv et al. (2009). The calculation assumes the *nominal* model of EBL in the range from 0.1 to 1000 μm (Aharonian 2001) and a typical gamma-ray power law photon distribution, $dn_\gamma/dE_\gamma \propto E_\gamma^{-2}$, in the range $10^3 \leq E_\gamma/m_ec^2 \leq 10^8$, for the blazar's spectrum.

Results for the pair beam generated according to the Monte Carlo model of the electromagnetic cascade are illustrate in Fig. 1. The top panel shows the peak (dash) and bulk (solid) energy of the pair beam, in units of

m_ec^2, as a function of distance from the blazar. The typical Lorentz factor of the pairs is a monotonically decreasing function of distance, in the range Γ=(5–1)$\times 10^5$ up to a Gpc away from the blazar. This shows that pair production typically peaks at near infrared-optical ($E_{\rm EBL} \simeq 0.1$–1 eV) EBL target photons interacting with gamma-rays with energy $E_\gamma \sim$ (0.1–1)TeV. The vertical bars indicate the energy range encompassing 68% of the particles. In fact, there is considerable energy spread about the mean value, which decreases towards larger distances as the energy range of gamma-ray photons interacting with the EBL is also reduced. However, the beam particles remain always ultra-relativistic, a detail relevant in the analysis below.

The bottom panel shows the angular distribution of beam pairs at distances of 55 Mpc (solid), 350 Mpc (black dash) and 1 Gpc (dot-dash) from the blazar. The angular distributions $dn/d\theta$ were simulated using analytical expressions from Schlickeiser et al. (2012) for angles in double photon collision. The beam angular spread is of order $\Delta\theta \simeq 10^{-5}$, consistent with Eq. (5) and the value of Γ estimated above. The colored, dash curves show the angular distribution of pairs at 350 Mpc (dash) in four different energy bins, indicating that the beam angular spread is primarily determined by low energy pairs with $\Gamma \simeq 10^4$–10^6 (red and yellow).

2.3. *IGM in Voids*

Free electrons in cosmic voids are characterized by a number density $n_{\rm v} \simeq 2\times 10^{-7}(1+\delta)(1+z)^3$ cm^{-3}, where the typical over-density is $\delta \lesssim -0.1$, and a temperature $T_{\rm v} \simeq$ a few $\times 10^3$ K (Peebles 1993).

3. BEAM INSTABILITY: REACTIVE VS KINETIC

The blazar induced pair beam is subject to microscopic instabilities, in particular two-stream like instabilities, of both electrostatic and electromagnetic nature. The beam is neutrally charged, so a return current is not induced. In the absence of magnetic field, $\omega_H \ll \omega_p$, with ω_H the cyclotron frequency, $\omega_p = (4\pi n_{\rm v} e^2/m_e)^{1/2}$ the plasma frequency of the IGM in voids and e the positron's charge, the instability is predominantly associated to Cherenkov emission of Langmuir waves, which operates under the following resonant condition

$$\omega - \mathbf{k}\cdot\mathbf{v} = 0, \tag{6}$$

where $\mathbf{k}$ is the wave-vector of the perturbation mode and $\mathbf{v}$ the beam particles velocity. The pair particles contribute equally to the dielectric function, as they have the same mass, number density, velocity distribution, and plasma frequency, $\omega_{p,b} = (4\pi n_b e^2/m_e)^{1/2}$. After separating the contributions from the background plasma and the beam particles, the dispersion relation for Langmuir waves, valid in the relativistic case, can be written as (Breizman 1990)

$$1 - \frac{\omega_p^2}{\omega^2} - \frac{4\pi e^2}{k^2}\int \frac{\mathbf{k}\cdot\partial f/\partial\mathbf{p}}{\mathbf{k}\cdot\mathbf{v}-\omega} d\mathbf{p} = 0. \tag{7}$$



FIG. 2.— Normalized growth rate, $\gamma_k/\pi\omega_p(n_b/n_v)$, from Eq. (11) in the plane $k_\parallel c/\omega_p - k_\perp c/\omega_p$. Bright is positive, dark is negative and the uniformly colored region is zero, as it lies outside the resonant region.

where, $f(\mathbf{p})$, is the distribution function of the beam particles. There are two important regimes that characterize the unstable behavior of the beam, namely reactive and kinetic. In the reactive case, the beam's velocity spread, $\Delta\mathbf{v}$, is negligible so all particles can participate to the unstable behavior and the growth rate of the instability is therefore fastest. In the kinetic regime, on the other hand, the velocity spread is considerable and only the resonant particles contribute to the growth of Langmuir waves, so the growth rate is slower than in the reactive case. Formally, the reactive regime is applicable when (Breizman & Ryutov 1971)

$$|\mathbf{k}\cdot\Delta\mathbf{v}| \ll \gamma_r, \tag{8}$$

where γ_r is the reactive growth rate. In this case, the integral in Eq. (7) can be solve in a simplified way, which involves neglect of the velocity spread around the mean value. This leads to the estimate of the reactive growth rate which, maximized along the longitudinal component of the wave-vector reads (Fainberg et al. 1970)

$$\gamma_r \simeq \omega_p \left(\frac{n_b}{\Gamma n_{\rm v}}\right)^{\frac{1}{3}} \left(\frac{k_\parallel^2}{k^2\Gamma^2} + \frac{k_\perp^2}{k^2}\right)^{\frac{1}{3}}, \tag{9}$$

with $k_\parallel = \omega_p/v$, and $k_\parallel$, $k_\perp$ the components of the wave-vector parallel and perpendicular to the beam direction, respectively. It is well known that, for an ultra-relativistic beam ($\Gamma \gg 1$), the fastest growing modes in the reactive regime are those quasi-perpendicular to the beam. This is due to the large suppression caused by relativistic inertia along the longitudinal direction (Fainberg et al. 1970). However, as shown later, for quasi-perpendicular directions of the wave vector, the reactive regime is not applicable.

When the approximation (8) is not valid, the growth rate is evaluated from a pole of the integrand in the dispersion relation (7), namely

$$\gamma_k = \omega_p \frac{2\pi e^2}{k^2}\int \mathbf{k}\cdot\frac{\partial f}{\partial\mathbf{p}}\,\delta(\omega_p - \mathbf{k}\cdot\mathbf{v})\,d\mathbf{p}. \tag{10}$$

If, as is the case here, despite the energy spread the particles remain ultra-relativistic and, $|\mathbf{v}| = c$,



FIG. 3.— Normalized growth rate as a function of $k_\parallel c/\omega_p$ inside the the resonant region, for values of $k_\perp$ where γ_k reaches its maximum values.



FIG. 4.— Normalized growth rate as a function of $k_\parallel c/\omega_p$ inside the the resonant region, for large values of $k_\perp$ where γ_k starts to drop compared to its maximum value.

can be assumed, the above integral can be simplified to (Breizman & Mirnov 1970)

$$\gamma_k = -\omega_p \pi \frac{n_b}{n}\left(\frac{\omega_p}{kc}\right)^3 \int_{\mu_-}^{\mu_+} d\mu \frac{2g + (\mu - \frac{k_\parallel c}{\omega_p})\frac{\partial g}{\partial \mu}}{[(\mu_+ - \mu)(\mu - \mu_-)]^{\frac{1}{2}}}. \quad (11)$$

Here, the integration variable μ is the angle between the particles and the beam direction, and

$$\mu_\pm = (\omega_p/kc)(k_\parallel/k \pm k_\perp/k\sqrt{k^2c^2/\omega_p^2 - 1}), \quad (12)$$

$$g(\theta) = \frac{m_e c}{n_b}\int pf(p,\theta)dp \simeq \frac{1}{\Gamma_{beam}\Delta\theta^2}e^{-\frac{\theta^2}{\Delta\theta^2}}. \quad (13)$$

The second equality for $g(\theta)$ in Eq. (13) is found to be a very good approximation based on results of the Monte Carlo model of the cascade. The integral for the growth rate in Eq. (11) can be evaluated numerically. The *qualitative* behavior of the growth rate, γ_k, on the plane $k_\parallel - k_\perp$ is summarized in Fig. 2 (Breizman & Ryutov 1971) for a beam at a Gpc from the blazar. Outside the narrow resonant region of k-space, corresponding in the plot to the uniform color, the growth rate is effectively null. In the narrow resonant region around $k_\parallel = \omega_p/c$, the growth can be positive (bright), negative (dark) and null, and for large enough values of $k_\perp$, it carries the sign of, $\omega_p/k_\parallel - c$ (see below). Within the resonant region, the growth rate as a function of $k_\parallel$ has typically two extrema, a maximum and a minimum. This is shown in Fig. 3 for values of $k_\perp$ of interest, i.e. where γ_k reaches its maximum values. The growth rate has its largest values where $k_\perp c/\omega_p \lesssim 1$, and decays rapidly in the opposite limit. This can be seen from Fig. 4 where γ_k is plotted for values of $k_\perp$ close to and much larger than ω_p/c. Finally, we find that the growth rate, maximized with respect to $k_\parallel$ and as a function of $k_\perp$, can be well approximated by the following expression by Breizman & Ryutov (1971)

$$\gamma_k \simeq \omega_p \frac{n_b}{\Gamma n_{\rm v}}\frac{1}{\Delta\theta^2}\frac{\omega_p^2}{\omega_p^2 + k_\perp^2 c^2}, \quad (14)$$

which we will be using in the following.

3.1. *Fastest Growing Modes*

For particles of an ultra-relativistic beam with modest angular spread, $\Delta\theta \ll 1$, we can assume $v_\parallel \simeq c$ and $v_\perp \simeq c\Delta\theta$. If the energy spread, $\Delta E/E \lesssim 1$, then the longitudinal velocity spread of the beam is

$$\Delta v_\parallel \simeq c\frac{\Delta E}{\Gamma^2 E} + c\Delta\theta^2. \quad (15)$$

Thus, for the angular spread and Lorentz factor characteristic of the blazars induced beam, the longitudinal velocity spread is negligible with respect to the perpendicular velocity spread. It turns out that, the first and second terms in Eq. (15) are comparable to within a factor of a few, so for the sake of simplicity in the following we retain the second term only, neglecting any fudge factor. We then combine Eq. (8), (9) and (15) to obtain the condition for applicability of the reactive regime, i.e.

$$2\Delta\theta \ll -\frac{k_\perp}{k_\parallel} + \left[\frac{k_\perp^2}{k_\parallel^2} + 4\left(\frac{n_b}{\Gamma n_{\rm v}}\right)^{\frac{1}{3}}\left(\frac{k_\parallel^2}{k^2\Gamma^2} + \frac{k_\perp^2}{k^2}\right)^{\frac{1}{3}}\right]^{\frac{1}{2}}. \quad (16)$$

This threshold value for the angular spread of the beam is plotted in Fig. 5 as a function of the wave-vector orientation, for different values of the Lorentz factor of the beam, and the beam-to-IGM electron density ratio. For given beam characteristics, the nature of the instability depends on the orientation of the wave-vector



FIG. 5.— Threshold value of the angular spread for kinetic regime, as a function of the wave-vector orientation for the case of a beam-to-IGM electron density ratio of 10^{-14} (red) and 10^{-15} (cyan), respectively, and for a beam Lorentz factors $\Gamma = 10^5$ (solid) and 5×10^5 (dash), respectively.

with respect to the beam and modes of reactive and kinetic character can appear simultaneously. In particular, modes close to perpendicular require a kinetic description. This is because the beam velocity spread along the wave-vector cannot be neglected.

Using again Eq. (8)–(15) with the estimate of the beam angular spread and bulk Lorentz factor from the previous Section, we find that the transition from reactive to kinetic regimes occurs for

$$\frac{k_\perp}{k} \simeq \times 10^{-5} \left(\frac{n_b/n_{\rm v}}{10^{-15}}\right) \left(\frac{\Gamma}{10^5}\right)^{-1} \left(\frac{\Delta\theta}{10^{-5}}\right)^{-3}. \qquad (17)$$

In the above estimate we have used a conservative value for the the ratio, $n_b/n_{\rm v}$. However, this number may can be much smaller due to distance from the blazars and to redshift evolution of the EBL, so that virtually the reactive regime may never apply. In any case, for perturbation modes more aligned to the beam direction than the above value, the instability is reactive in nature, but the growth rate decreases due to the increasing relativistic inertia. In the opposite case, of larger $k_\perp/k$, the instability is kinetic and the growth rate also decreases because of the increasing velocity spread along $\mathbf{k}$. Thus, perturbations with $k_\perp/k$ roughly as in Eq. (17) grow at the fastest rate, given by

$$\gamma_{max} \simeq \omega_p \frac{n_b}{\Gamma n_{\rm v}} \frac{1}{\Delta\theta^2_{beam}}, \qquad (18)$$

which can be actually derived either from the above reactive or kinetic expressions for the growth rate.

4. BEAM STABILIZATION

The generation of Langmuir waves induces particle-wave interactions which cause the particles to diffuse in momentum space, eventually leading to the relaxation of the beam (Melrose 1989). In the very first phase of relaxation, described by the quasilinear approximation, the generation of Langmuir waves is balanced by their drift at the group velocity, $v_g \simeq 3v_t^2/c$, to regions of lower wave energy density. However, due to the smallness of the group velocity and to the large scale of the wave energy gradient, of order the pair beam distance to the blazar, the timescale associated to this process largely exceeds a Hubble time. Therefore in the quasilinear relaxation the growth of the Langmuir waves is uncompensated.

4.1. *Nonlinear Landau damping*

The main process that compensates the growth of Langmuir waves is induced scattering off plasma ions, also known as non-linear Landau damping (Tsytovich & Shapiro 1965; Breizman et al. 1971; Lesch & Schlickeiser 1987). In this process, a thermal ion, with characteristic velocity, v_{ti}, interacts with the beat wave produced by two Langmuir oscillations, $\omega(\vec{k})$, $\omega(\vec{k}')$, under the condition for Cherenkov interaction, i.e.

$$\omega(\vec{k}) - \omega(\vec{k}') = (\vec{k} - \vec{k}') \cdot \vec{v}_{ti}. \qquad (19)$$

In a thermal plasma, the scattering process leads to damping of the beat wave and a reduction of its frequency (Kaplan & Tsytovich 1973). Due to their dispersion relation, Langmuir waves are consequently shifted towards regions of phase-space of lower wave-vectors. Eventually their phase speed exceeds the speed of light, $\omega/k > c$, and they fall out of resonance with the beam particles. The change in wave-vector suffered in one scattering process is of order (Kaplan & Tsytovich 1973)

$$\Delta k \sim \omega_p \frac{v_{ti}}{v_{te}} = 50 \times \frac{\omega_p}{c} \left(\frac{T_{\rm v}}{10^3 {\rm K}}\right)^{-\frac{1}{2}}, \qquad (20)$$

where v_{te} is the electron thermal speed. Since for Langmuir waves, $k \sim \omega_p/c$, the above equation implies $\Delta k/k \gg 1$, i.e. at the low temperatures characteristic of the IGM in voids, waves are kicked out of resonance in a single scattering event. The rate of induced scattering off thermal ions is (Breizman et al. 1971)

$$\gamma_{\rm nl} = \omega_p \frac{W}{n_{\rm v} k_B T_{\rm v}}, \qquad (21)$$

where W is the energy density of Langmuir waves. The latter evolves in time according to

$$\frac{\partial W}{\partial t} = 2(\gamma_{max} - \gamma_{\rm nl})W. \qquad (22)$$

Over time, a quasi steady-state condition will be established such that the growth of waves due to the beam instability is balanced on average by induced scattering. By equating the two competing rates we can obtain an estimate for the equilibrium value of W. Expressed in terms of the energy density of the beam, the latter is

$$\frac{W_{eq}}{n_b \Gamma m_e c^2} \simeq 10^{-8} \left(\frac{T_{\rm v}}{10^3 {\rm K}}\right) \left(\frac{\Gamma}{3\times10^5}\right)^{-2} \left(\frac{\Delta\theta}{10^{-5}}\right)^{-2}.$$

The power associated with the emission of Langmuir waves by the beam is then $dW/dt = 2W_{eq}\gamma_{max}$. The corresponding beam relaxation timescale is

$$\tau_{beam} = \frac{n_b \Gamma m_e c^2}{2 W_{eq} \gamma_{max}} = 10^{12}\mathrm{yr} \left(\frac{\Gamma}{10^5}\right)^4 \times \left(\frac{\Delta\theta}{10^{-5}}\right)^4 \left(\frac{D}{\mathrm{Gpc}}\right)^2 \left(\frac{T_{\rm v}}{10^3\mathrm{K}}\right)^{-1} (1+z)^{7-\zeta} \qquad (23)$$

where redshift dependence is made explicit and $\delta = -0.7$ has been used. The above timescale is generally longer than the Hubble time, and even at close distances from the blazar, remains longer than the pairs cooling time on the Cosmic Microwave Background, $\ell_{IC}/c \simeq \times 10^6 (E_\pm/TeV)^{-1}(1+z)^{-4}$ yr. This suggests that the beam instability plays only a secondary role on the electromagnetic shower and the beam dynamics and, as a consequence, the thermal history of the IGM and/or structure formation.

4.2. *Plasma Inhomogeneities*

In addition to induced scattering by thermal ions, plasma inhomogeneities can also induce loss of resonance between particles and Langmuir waves (Breizman & Ryutov 1971; Nishikawa & Ryutov 1976). The rate of change of the wave-vector due to inhomogeneities is given by $d\vec{k}/dt = -\vec{\nabla}\omega$. For Langmuir waves the RHS of the above equation is mostly contributed to by density inhomogeneities. In the following we consider only the effects of inhomogeneities on the longitudinal component of the wave-vector. These are described by the equation

$$\frac{dk_\parallel}{dt} \simeq \frac{1}{2}\frac{\omega_p}{\lambda_\parallel}, \qquad (24)$$

with $\lambda_\parallel = n_{\rm v}/(\vec{\nabla} n_{\rm v})_\parallel$, the length scale of the density gradient along the beam.

Fig. 6 shows the range of scale lengths of IGM density gradients as a function of IGM gas over-density, for three different cosmological redshifts, $z = 0$ (top), $z = 1$ (middle) and $z = 3$ (bottom). The data is extracted from a cosmological simulation of structure formation including hydrodynamics, dark matter, and self-gravity as described in Miniati & Colella (2007). A flat ΛCDM universe is adopted with the following cosmological parameters: total mass density, normalized to the critical value for closure, $\Omega_m = 0.2792$, normalized baryonic mass density, $\Omega_b = 0.0462$, normalized vacuum energy density, $\Omega_\Lambda = 1 - \Omega_m = 0.7208$, Hubble constant $H_0 = 70.1$ km s^{-1} Mpc^{-1}, spectral index of primordial perturbation, $n_s = 0.96$, and rms linear density fluctuation within a sphere with a comoving radius of 8 h^{-1} Mpc, $\sigma_8 = 0.817$, where $h \equiv H_0/100$ (Komatsu et al. 2009). The computational box has a comoving size $L = 50h^{-1}$ Mpc, is discretized with 512^3 comoving cells, corresponding to a nominal spatial resolution of $100h^{-1}$ comoving kpc. The collisionless dark matter component is represented with 512^3 particles with mass $6 \times 10^5 h^{-1}$ M$_\odot$.

According to Fig. 6, the distance covered by the beam particles during the fastest growth time, $\sim c\gamma_{max}^{-1} \simeq 1$ kpc, is much shorter than the typical scale-length of density gradients. This correspond to the case of regular, as opposed to random, inhomogeneities. In this case, the condition for wave excitation can be expressed as (Breizman & Ryutov 1971)



FIG. 6.— Characteristic length scale of density gradient in the IGM as a function IGM gas over-density. The shaded area extends ± one root-mean-squared value about the average.

$$\gamma_{max} \frac{\Delta k_\parallel}{|dk_\parallel/dt|} > \Lambda_c, \qquad (25)$$

where Λ_c is the Coulomb logarithm and $\Delta k_\parallel$ the change in longitudinal component of the wave-vector allowed by the resonant condition (6). Eq. (25) states that, in order for the wave energy to grow, the resonance condition must be violated only after several growth times have lapsed. Using Eq. (6) (and neglecting the term $\Delta E/E\Gamma^2$) to obtain, $\Delta k_\parallel \lesssim \frac{\omega_p}{c}\Delta\theta^2 + k_\perp \Delta\theta$, Eq. (25) can be solved to express the condition for wave excitation in terms of $\lambda_\parallel$, i.e.

$$\lambda_\parallel \geq \frac{c}{2\omega_p}\frac{\Gamma n_{\rm v}}{n_b}\Lambda_c \left(1 + \frac{k_\perp}{k_\parallel \Delta\theta}\right)^{-1} \simeq 1.5 \times 10^7 \mathrm{kpc}\, \Lambda_c \left(\frac{\Gamma}{10^5}\right)\left(\frac{n_{\rm v}/n_b}{10^{15}}\right), \qquad (26)$$

where we have used, $\omega_p = 16 s^{-1}$ for the plasma frequency in voids. The above estimate gives an extremely stringent constraint on the minimum allowed scale-length of density gradients required for the growth of Langmuir waves, which is orders of magnitude larger than the values reported in Fig. (6), even when using a very conservative value for the ratio, $n_{\rm v}/n_b$. Therefore, such constraints cannot be possibly satisfied due to the presence of matter fluctuations associated to cosmic structure, providing another independent argument against the growth of Langmuir waves and the unstable behavior of the pair beam.

5. CONCLUSION

We considered the stability properties of a low density ultra relativistic pair beam produced in the intergalactic medium by multi-TeV gamma-ray photons from blazars. The physical properties of the pair beam are determined through a Monte Carlo model of the electromagnetic cascade. In summary we find that the combination of kinetic effects, non-linear Landau damping and density inhomogeneities appear to stabilize blazars induced ultra-relativistic beams over a Hubble timescale, so that the electromagnetic cascade remains unaffected by the beam instability. This implies that the lack of a bumpy feature at multi-GeV energies in the gamma-ray spectrum of distant blazars cannot be attributed such instabilities. Finally, heating of the IGM by pair beams appears completely negligible.

Useful exchange and comments by B. N. Breizman, D. D. Ryutov and R. Schlikeiser are gratefully acknowledged. We are grateful to D. Potter for making available is `grafic++` package for cosmological initial conditions. The numerical calculations were per performed at the Swiss National Supercomputing Centre.

REFERENCES

Aharonian, A. F., arXiv:astro-ph/0112314 (2001)

Aharonian, F. A. et al. 2006, Nature, 440, 1018

Benz, A. 1993, Plasma Astrophysics (Kuwler: Dordrecht)

Bertschinger, E. 2001, ApJS, 137, 1

Bonometto, S., Rees, M. R. 1971, MNRAS, 152, 21

Breizman, B. N. 1990, Reviews of Plasma Physics, 15, 61 (ed. B. B. Kadomstev)

Breizman, B. N., Mirnov, V. V. 1970, Geomagn. Aeron., 10, 34

Breizman, B. N., Ryutov, D. D. 1971, JETP, 33, 220

Breizman, B. N., Ryutov, D., Chebotaev, P. Z. 1972, JETP, 35, 72

Breizman, B. N., Ryutov, D. D. 1974, Nucl. Fusion, 14, 873

Broderick, A. E., Chang, P., Pfrommer, C. 2012, ApJ, 752, 22

Chang, P., Broderick, A. E., Pfrommer, C. 2012, ApJ, 752, 23

Dermer, C. D., Cavadini, M., Razzaque, S., Finke, J. D., Chiang, J., Lott, B. 2011, ApJ, 733, L21

Elyiv, A., Neronov, A., D. V. Semikoz, D. V. 2009, Phys. Rev. D, 80, 023010

Ensslin, T. A., Pfrommer, C., Miniati, F., Subramanian, K. 2011, A&A, 527, A99

Fainberg, Ya. B., Shapiro, V. D., Shevchenko, V. I. 1970, JETP, 30, 528

Franceschini, A., Rodighiero, G., Vaccari, M. 2008, A&A, 487, 837

Gould, R. J. & Schréder, G. P. 1967, Phys. Rev. Lett., 16, 252

Kaplan, S. A., Tsytovich, V. N. 1973, Plasma Astrophysics, (Pergamon Press: Oxford)

Kneiske, T. M., Bretz, T., Mannheim, K., Hartmann, D. H. 2004, A&A, 413, 807

Komatsu, E., et al. 2009, ApJS, 180, 330

Lesch, H., Schlickeiser, R. 1987, A&A, 179, 93

Melrose, D. B. 1989, Instabilities in Space and Laboratory Plasmas (Cambridge University Press)

Miniati, F., Bell, A. R., 2011, ApJ, 729, 73

Miniati, F., & Colella, P. 2007, Journal of Computational Physics, 227, 400

Neronov, A., Semikoz, D. 2009, Phys. Rev. D, 80, 123012

Neronov, A., Vovk, I. 2010, Science, 328, 7

Nishikawa, K, Ryutov, D. D. 1976, Journal of the Physical Society of Japan, 41, 1757

Peebles, P. J. E. 1993, Principles of Physical Cosmology (Princeton New Jersey: Princeton University Press)

Pfrommer, C., Chang, P., Broderick, A. E. 2012, ApJ, 752, 24

Plaga, R. 1995, Nature, 374, 430

Schlickeiser, R., Elyiv, A., Ibscher, D., Miniati, F. 2012, ApJ, submitted

Tavecchio, F., Ghisellini, G., Bonnoli, G., Foschini, L. 2011, MNRAS, 414, 3566

Taylor, A. M., Vovk, I., Neronov, A. 2011, A&A, 529, A144

Tsytovich, V. N., Shapiro, V. D. 1965, Nuclear Fusion, 5 228